- MEYRICK, E. (1928) A Revised Handbook of British Lepidoptera, pp. 222-223. Watkins & Doncaster, London.
- NEWMAN, E. (1869) The Natural History of British Moths, pp. 136-138. Tweedie, London.
- PETERSEN, W. (1909) Ein Beitrag zur kenntnis der Gattung Eupithecia. Deutsche Zeitschrift "Iris", 22: 242.
- PIERCE, F. N. (1911) The Genitalia of the Geometridae, p.49. Northern Publishing Company, Liverpool.
- SKINNER, B. (1984) Colour Identification Guide to the Moths of the British Isles, p.41. Viking, London.
- SOUTH, R. (1961) The Moths of the British Isles, 2: 218-219. Warne, London.
- STOKOE, W. J. (1948) The Caterpillars of the British Moths, 2: 129-130. Warne, London.
- TUTT, J. W. (1899) The Natural History of the British Lepidoptera, 1: 85-86. Swan Sonnenschein, London.
- TUTT, J. W. (1906) Practical Hints Relating to the Eupitheciids. Entomologist's Rec. J. Var., 18: 219.

NOTE: in an interesting paper (Fibiger, M. Lepidoptera N. S. III 307-311 (1980) and Kaaber, S. Lepidoptera N. S. IV41-29 (1982)) the authors consider the status of the Eupithecia couples absinthiata/ goossensiata and innotata/fraxinata. Their major conclusion on the former pair, based on morphology, genitalia examination and captive breeding is that E. goossensiata is but a race or subspecies of absinthiata. In Denmark goossensiata is widely distributed on sandy soil, whereas absinthiata is mainly restricted to woodland habitats on clay soil. PAS.

OCHTHERA SPP. (DIPT.: EPHYDRIDAE): A CORRECTION. – In 1983, Ent. Rec. 95: 154 I reported a specimen of the uncommon fly Ochthera mantis Deg. from Catfield Fen, Norfolk, remarking that I had no knowledge of its distribution. Since that was written, however, Dr. A. G. Irwin has published an important paper in which he distinguishes a third British species of the genus, Ochthera manicata F., and shows that all captures in E. Anglia known to him, centring on this small area of N. E. Norfolk, are referable to O. manicata (1985, Ent. mon. Mag. 121: 151-4). It therefore is no surprise that my Catfield specimen is now found to belong to the last-named species and not O. mantis; particularly as it has occurred at the same locality to both Dr. Irwin and also Dr. J. W. Ismay (Irwin, *l.c.*). O. mantis on the other hand, though thinly scattered over the British Isles, is not hitherto known from that region. – A. A. ALLEN.